



ClenbuGen 40 (Clenbuterol)

40 mcg/each

Oral Tablets

FORMULA

Each uncoated tablet contains: Clenbuterol hydrochloride (40 mcg), Excipients (q.s.)

DOSAGE AND USE

Human Warnings: Not for human use. Do not use in horses intended for human consumption. Keep out of the reach of children. In case of accidental ingestion, contact a physician immediately. Ingestion of Ventipulmin Syrup may cause undesirable reactions. Clenbuterol, like other beta adrenergic agonists, can produce significant cardiovascular effects in some people as evidenced by elevated pulse rate, blood pressure changes and/or ECG changes.

Initial dosage: administer 0.5 mL/100 lbs (0.8 mcg/kg) for 3 days (6 treatments).

CHARACTERISTICS AND INDICATIONS

ClenbuGen 40 product with bronchial-dilation action. It relaxes smooth muscles of the bronchi, improving their road, relieves tension in the chest, wheezing and eases breathing. It is used for removal of bronchospasm in asthma and other pulmonary diseases.

ADVERSE REACTIONS

Pregnancy and lactation: Ask your doctor or pharmacist before taking any medicine. ClenbuGen 40 is not applied in the first three months of pregnancy, and the latter because of its depressing effect on parturition. ClenbuGen 40 is not recommended in breastfeeding women.

Driving and using machines: The ClenbuGen 40 can cause hand tremors and restlessness, which requires caution when driving or operating machines during treatment with it.

CONTRAINDICATIONS

Do not take ClenbuGen 40 at : allergy (svara chuvstvitelnost) to the active substance or to any of the ingredients of the product; thyrotoxicosis (overactive thyroid gland); cardiovascular disease - ischemic heart disease (poor circulation to the heart muscle), hypertension, hypertrophic obstructive cardiomyopathy, and especially those associated with abnormal heart rhythm (tachyarrhythmia); recent myocardial infarction; first trimester of pregnancy.

Important information about some of the ingredients of ClenbuGen

- The medicinal product contains lactose excipient. If you have an intolerance to some sugars, consult your doctor before taking this medicine.
- The composition of the product is included wheat starch. Not pose a risk for people with celiac disease. Patients with wheat allergy (different from celiac disease) should not take this medicine.

DRUG INTERACTIONS

You should not take concomitantly with ClenbuGen 40 :

other medications for the treatment of bronchospasm (e.g. salbutamol, theophylline, etc. nov phyllin) corticosteroids, water tablets (diuretics), drugs for treatment of heart failure (digitalis), halothane and other means of anesthesia, drugs for the treatment of depression (MAO- inhibitors or tricyclic antidepressants), drugs for treating diseases of the thyroid (thyroid hormone)

- can increase the risk of adverse effects; drugs for the treatment of diabetes
- ClenbuGen 40 may reduce their therapeutic effect; beta-blockers (medicines used to treat hypertension eg. propranolol, atenolol, metoprolol) and ClenbuGen 40 antagonize their interaction effects during coadministration.

OVERDOSE

In case of adverse effects (shaky hands, sweating, headache, palpitations...) withdraw use of the drug and immediately call your doctor or pharmacist.

PACKAGING

Box with 50 tablets of 40 mcg

STORAGE CONDITIONS

Store in a cool dry place below 30°C.

Protect from light. Keep out of reach of children.

Sale only with Prescription.

Made in India