



# StanoGen 10 (Stanozolol)

**10 mg/each**

**Oral Tablets**

## FORMULA

Each uncoated tablet contains: Stanozolol USP (10 mg), Excipients (q.s.)

## DOSAGE AND USE

Usual Adult Dose for angioedema: Prophylactic use to decrease frequency and severity of attacks of hereditary angioedema.

Initial dose: 2 mg orally 3 times a day.

## THERAPEUTIC ACTION

Anabolic hormone that promotes the processes of reconstruction of body tissues and reverses body catabolism. It is used therapeutically to treat hereditary angioedema, the cryofibrinogenemia decreasing the severity and frequency of seizures.

## MECHANISM OF ACTION AND PHARMACOKINETICS

Stanozolol is a synthetic steroid derived from testosterone that enhances muscular development. However, it also has the androgenic properties of the sexual male hormones (testosterone). Stanozolol increases plasma levels of the LDL and reduces the levels of HDL, even though total cholesterol or triglycerides are not modified. When administration is discontinued, this levels turn back to normal. Elimination is through the kidneys, a small part is eliminated through feces, because of the enterohepatic incubation.

## CONTRAINDICATIONS

Use of stanozolol is contraindicated in male patients with breast carcinoma or with suspected or latent prostate carcinoma. In women, stanozolol is contraindicated in breast cancer with hypercalcemia. Androgenic steroids can stimulate the osteolytic bone resorption.

Stanozolol should be used with caution in children and adolescents because it can cause an early closure of the epiphysis, precocious sexual development in males and masculinization in female. Epiphysis maturation may be faster than the growth and continue 6 months after discontinuation of the medication.

## PRECAUTIONS AND WARNINGS

Anabolic steroids should be used with caution in children and adolescents because it can cause an early closure of the epiphysis, precocious sexual development in males and masculinization in female. Epiphysis maturation may be faster than the growth and continue 6 months after discontinuation of the medication.

## DRUG INTERACTIONS

Effects of heparin and oral anticoagulants may be increased by androgens, including stanozolol.

Antidiabetics (insulin): there are studies in which a potentiation of the hypoglycemic effect has been shown, with toxicity risk. The mechanism has not been established, but it has been suggested that with the increase of metabolic activity by anabolics and increasing the glucose consumption, the effect of the hypoglycemic can be increased.

## RESTRICTIONS IN USE

Do not use in cases of hypersensitivity to any of the components, pregnancy or lactancy.

## OVERDOSE

In case of adverse effects withdraw use of the drug and immediately call your doctor or pharmacist.

## PACKAGING

Box with 50 tablets of 10 mg

## STORAGE CONDITIONS

Store in a cool dry place below 30°C.

Protect from light. Keep out of reach of children.

**Sale only with Prescription.**

**Made in India**