



TrenaGen Mix 150 (Trenbolone Compound Injection)

For I.M. use only

FORMULA

Each ampoule of 1 ml contains: Trenbolone Acetate USP (50 mg), Trenbolone Enanthate (50 mg), Trenbolone Hexahydrobenzylcarbonate (50 mg), Oily base (q.s.)

ADMINISTRATION

Intramuscular injection is to be administered deep into the muscle, due to its oily vehicle. Do not administer intravenously. Shake well before use. Any adverse effect after the use of this product please report in writing to the National Health Surveillance India.

This drug should be inspected visually for particulate matter and discoloration prior to administration. Warming and shaking the vial should dissolve any crystals that may have formed during storage at temperatures lower than recommended.

INDICATIONS AND USAGE

Males: TrenaGen Mix 150 use may be indicated in patients where substantial weight gain and increases in musculature are required for patient health after substantial losses of body mass especially in instances where caloric intake is limited and other anabolic therapies have previously failed. The physician and patient must consider the risks of therapy versus the potential benefits.

Active life: 8 days

Detection time: up to 5 months

Anabolic/Androgenic ratio: 500/500

SIDE EFFECTS

Male: Gynecomastia, excessive frequency and duration of penile erections, oligospermia. Skin and Appendages: Hirsutism, male pattern baldness and acne, gynecomastia. Fluid/electrolyte Disturbances: Retention of sodium, chloride, water, potassium, calcium, and inorganic phosphates.

Gastrointestinal: Nausea, cholestatic jaundice, alterations in liver function tests; hepatocellular neoplasms, peliosis hepatis, hepatic adenomas, nephritis, and cholestatic hepatitis.

Hematologic: Suppression of clotting factors II, V, VII, & X; bleeding in patients on anticoagulant therapy.

Neurological: Increased libido, headache, anxiety, depression, extreme agitation, irritability, and generalized paresthesia.

TrenaGen Mix 150 may cause severe aggressive behavior.

OVERDOSE

In case of adverse effects withdraw use of the drug and immediately call your doctor or pharmacist.

PACKAGING

Box with 5 ampoules of 1 ml.

STORAGE CONDITIONS

Store in a cool dry place below 30°C. Do not freeze.

Protect from light. Keep out of reach of children.

Sale only with Prescription.

Made in India