



# Turigen 10 (Chlorodehydromethyltestosterone)

**10 mg/each**  
Oral Tablets

## FORMULA

Each uncoated tablet contains: Chlorodehydromethyltestosterone (10 mg), Excipients (q.s.)

## CHARACTERISTICS AND INDICATIONS

TuriGen is characterized for having anabolic and androgenic properties, being the main indication of these the treatment of male hypogonadism.

It enhances protein synthesis because it supports protein increase at muscular level. This effect is shown in a positive balance of nitrogen and a feeling of improved wellbeing. Calcium balance is also positively influenced since it promotes the calcium deposit in bones. TuriGen is indicated in the treatment of all diseases and conditions in which an anabolic effect is needed (increase of protein synthesis) and a general strengthening effect.

## ADVERSE REACTIONS

The following reactions require medical supervision and are common with the use of all types of testosterone:

Menstrual irregularities in women, bladder irritability, gynecomastia, anaphylaxis, oedema, erythrocytosis, gastrointestinal irritation, hypercalcemia and polycythemia, androgenic alopecia, seborrhea and acne in women and men.

Prostatic hypertrophy Carcinoma, and increase of sexual desire, cholestatic jaundice, excitability; hypercholesterolemia; hypercalcemia, oedema.

In women: virilization

In men: priapism, oligospermia, hirsutism, decreased libido, decreased testicular mass, functional impotence.

Other adverse reactions are: constipation, nausea, diarrhea, infection, changes in libido, stomach ache, difficulty sleeping, headache, anxiety, depression, generalized paresthesia, rash.

## CONTRAINDICATIONS

Hypersensitivity to Methandrostenolone.

Severe hepatic impairment.

In prepubertal individuals or individuals in aggressive state.

Pregnancy, breastfeeding.

## DRUG INTERACTIONS

As with testosterone and other androgens and anabolic steroids they increase the activity of certain drugs and consequently its toxicity. Among them are cyclosporine, antidiabetics, levothyroxine, anticoagulants such as warfarin.

A resistance to neuromuscular blockers has also been reported.

## PRECAUTIONS AND WARNINGS

TuriGen is a  $17\alpha$ -alkylated substance associated with hepatotoxicity. Hepatic function should be closely monitored during treatment. It should not be administered to patients with hepatic impairment.

## OVERDOSE

In case of adverse effects withdraw use of the drug and immediately call your doctor or pharmacist. Support measures must be taken in case of an overdose, such as keeping an open airway and monitoring cardiovascular function.

## PACKAGING

Box with 50 tablets of 10 mg

## STORAGE CONDITIONS

Store in a cool dry place below  $30^{\circ}\text{C}$ .

Protect from light. Keep out of reach of children.

**Sale only with Prescription.**

**Made in India**